


Spectroscopy results from the Tevatron



Heavy Quark and Leptons 2010 - Frascati, Oct 11, 2010

Diego Tonelli - Fermilab

The beginning, 33 years ago...



VOLUME 39, NUMBER 5

1 AUGUST 1977

the gauge is fixed up to bound
above results are encouraging.
that direct closed loop calculation

her, simply because dimensional
respects the gauge invariances)
tadpole diagrams.

Observation of a Dimuon Resonance at 9.5 GeV in 400-GeV Proton-Nucleus Collisions

S. W. Herb, D. C. Hom, L. M. Lederman, J. C. Sens,^(a) H. D. Snyder, and J. K. Yoh
Columbia University, New York, New York 10027

and

J. A. Appel, B. C. Brown, C. N. Brown, W. R. Innes, K. Ueno, and T. Yamanouchi
Fermi National Accelerator Laboratory, Batavia, Illinois 60510

and

A. S. Ito, H. Jöstlein, D. M. Kaplan, and R. D. Kephart
State University of New York at Stony Brook, Stony Brook, New York 11974

(Received 1 July 1977)

Accepted without review at the request of Edwin L. Goldwasser under policy announced 26 April 1976

Dimuon production is studied in 400-GeV proton-nucleus collisions. A strong enhancement is observed at 9.5 GeV mass in a sample of 9000 dimuon events with a mass $m_{\mu^+\mu^-} > 5$ GeV.

The Physics

QCD – a pristine example of successful QFT

However low-energy regime far from being understood. One of the few SM soft-spots remaining.



Major show-stopper in precision CKM predictions. Also affects hadron spectroscopy, decay rates, lifetimes.

Heavy quarks help. Static source of color field for light-quark partners

Some quantities calculable with lattice. Powerful numerical technique, not truly illuminating on underlying physics.

Effective theories – work-around that decouples dynamic degrees of freedom and provide quantitative predictions. Still pretty much disconnected from general theory.

Experimental input crucial to adjust inputs/approximations and discriminate among models. One day will hopefully provide clear-cut link with high-Q perturbative regime.


The ingredients

Large production x-sections of all HF in 1.96 TeV $ppbar$ collisions.

Tracking


$$\sigma_{p_T} / p_T^2 \sim 0.1\% / \text{GeV}$$

30 microns vertex resolution



Trigger

Online silicon tracking exploits long-lived fully-hadronic HF decays by triggering on displaced tracks



48 micron resolution impact par.




















Mu acceptance

$$|\eta| < 3, p_T > 1.5\text{-}2.0 \text{ GeV}$$

Tenths of millions J/ψ collected (20% from B)



The roadmap

	D ⁺ -D _s ⁺ mass difference	(2003)
 	X(3872) confirmation	(2004)
	B hadron masses	(2006)
 	Mass of exclusive B _c	(2006 and 2008)
	Orbitally excited D	(2006)
	X(3872) quantum num.	(2006 and 2007)
	Search for S=-2 pentaquarks	(2007)
	Σ _b observation	(2007)
 	Cascade b observation	(2007)
 	Orbitally excited B _s	(2008)
 	Orbitally excited B ⁰	(2009)
	Observation of Y(4140)	(2009)
 	Observation of Ω _b	(2008 and 2009)

DØ: so far 6 papers published:
2 **topcite50+** and 1 **topcite100+**

CDF: so far 17 papers published:
7 **topcite50+** and 1 **topcite100+**

The impact

PRL 102, 242002 (2009) PHYSICAL REVIEW LETTERS week ending 19 JUNE 2009

Evidence for a Narrow Near-Threshold Structure in the $J/\psi \phi$ Mass Spectrum in $B^+ \rightarrow J/\psi \phi K^+$ Decays

VOLUME 93, NUMBER 7 PHYSICAL REVIEW LETTERS week ending 13 AUGUST 2004

Observation of the Narrow State $X(3872) \rightarrow J/\psi \pi^+ \pi^-$ in $\bar{p} p$ Collisions at $\sqrt{s} = 1.96$ TeV

VOLUME 93, NUMBER 16 PHYSICAL REVIEW LETTERS week ending 15 OCTOBER 2004

Observation and Properties of the $X(3872)$ Decaying to $J/\psi \pi^+ \pi^-$ in $p \bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV

PRL 96, 082002 (2006) PHYSICAL REVIEW LETTERS week ending 3 MARCH 2006

Evidence for the Exclusive Decay $B_c^\pm \rightarrow J/\psi \pi^\pm$ and Measurement of the Mass of the B_c^\pm Meson

PRL 96, 102002 (2006) PHYSICAL REVIEW LETTERS week ending 17 MARCH 2006

Measurement of the Dipion Mass Spectrum in $X(3872) \rightarrow J/\psi \pi^+ \pi^-$ Decays

PRL 96, 202001 (2006) PHYSICAL REVIEW LETTERS week ending 26 MAY 2006

PRL 98, 132002 (2007) PHYSICAL REVIEW LETTERS week ending 30 MARCH 2007

Analysis of the Quantum Numbers J^{PC} of the $X(3872)$ Particle

PRL 99, 052001 (2007) PHYSICAL REVIEW LETTERS week ending 3 AUGUST 2007

Direct Observation of the Strange b Baryon Ξ_b^-

PRL 99, 052002 (2007) PHYSICAL REVIEW LETTERS week ending 3 AUGUST 2007

Observation and Mass Measurement of the Baryon Ξ_b^-

PRL 99, 172001 (2007) PHYSICAL REVIEW LETTERS week ending 26 OCTOBER 2007

Properties of $L = 1$ B_1 and B_2^* Mesons

PRL 100, 082001 (2008) PHYSICAL REVIEW LETTERS week ending 29 FEBRUARY 2008

Observation of Orbitally Excited B_s Mesons

PRL 100, 082002 (2008) PHYSICAL REVIEW LETTERS week ending 29 FEBRUARY 2008

Observation and Properties of the Orbitally Excited B_{s2}^* Meson

PRL 100, 182002 (2008) PHYSICAL REVIEW LETTERS week ending 9 MAY 2008

Observation of the Decay $B_c^\pm \rightarrow J/\psi \pi^\pm$ and Measurement of the B_c^\pm Mass

PRL 101, 012001 (2008) PHYSICAL REVIEW LETTERS week ending 4 JULY 2008

Observation of the B_c Meson in the Exclusive Decay $B_c \rightarrow J/\psi \pi$

PRL 101, 232002 (2008) PHYSICAL REVIEW LETTERS week ending 5 DECEMBER 2008

Observation of the Doubly Strange b Baryon Ω_b^-

PRL 102, 102003 (2009) PHYSICAL REVIEW LETTERS week ending 13 MARCH 2009

Measurement of Resonance Parameters of Orbitally Excited Narrow B^0 Mesons

PRL 103, 152001 (2009) PHYSICAL REVIEW LETTERS week ending 9 OCTOBER 2009

Precision Measurement of the $X(3872)$ Mass in $J/\psi \pi^+ \pi^-$ Decays

PHYSICAL REVIEW D 80, 072003 (2009)

Observation of the Ω_b^- baryon and measurement of the properties of the Ξ_b^- and Ω_b^- baryons

PRL 99, 202001 (2007) PHYSICAL REVIEW LETTERS week ending 16 NOVEMBER 2007

Observation of the Heavy Baryons Σ_b and Σ_b^*

PHYSICAL REVIEW D 73, 051104 (2006)



Measurement of mass and width of the excited charmed meson states D_1^0 and D_2^{*0} at CDF

ICHEP2010

Improved determination of $\Sigma_b^{()}$ b -baryons
resonance parameters*

$\Sigma_b^{(*)}$ - Intro and status

2006 – Evidence for new bottom baryons using 1 fb⁻¹



Update to 6 fb⁻¹, provide independent determination of masses, widths and isospin splittings

ICHEP2010


$\Sigma_b^{(*)}$ – selection & reconstruction

5-track final states.

Associate soft pion to large and clean signal of all-hadronic Λ_b decays:

$$\Sigma_b^{\pm(*)} \rightarrow \Lambda_b \pi^{\pm}$$

Mass-difference to cancel Λ_b mass resolution and several systematic uncertainties.





$$Q = M(\Sigma_b^{(*)} \rightarrow \Lambda_b^0 \pi_{\text{soft}}^{\pm}) - M(\Lambda_b^0) - m(\pi^{\pm})_{PDG}$$

- ❑ Long Λ_b lifetime,
- ❑ Large momentum of Λ_b pion,
- ❑ Prompt production of pion from Σ

ICHEP2010

$\Sigma_b^{(*)}$ – mass difference fit





Non-relativistic P-wave BW convoluted with resolution for signal. Empirical background. All signals comfortably beyond 5 σ significance

ICHEP2010

$\Sigma_b^{(*)}$ – Systematic uncertainties

Dominated by tracking resolution of soft pion from Σ_b . Compare data and simulation for large samples of $D^* \rightarrow D^0 \pi$ decays.



25% relative uncertainty.

Minor contributions from magnetic field knowledge, Λ_b mass uncertainty, alternate background models.

ICHEP2010

 $\Sigma_b^{(*)}$ – final results

Using CDF measurement of $5619.7 \pm 1.2 \pm 1.2$ MeV for the Λ_b mass

State	$M, \text{ MeV}/c^2$	$\Gamma_0, \text{ MeV}/c^2$
Σ_b^+	$5811.2^{+0.9}_{-0.8} (\text{stat}) \pm 1.7 (\text{syst})$	$9.2^{+3.8}_{-2.9} (\text{stat})^{+1.0}_{-1.1} (\text{syst})$
Σ_b^{*+}	$5832.0 \pm 0.7 (\text{stat}) \pm 1.8 (\text{syst})$	$10.4^{+2.7}_{-2.2} (\text{stat})^{+0.8}_{-1.2} (\text{syst})$
Σ_b^-	$5815.5^{+0.6}_{-0.5} (\text{stat}) \pm 1.7 (\text{syst})$	$4.3^{+3.1}_{-2.1} (\text{stat})^{+1.0}_{-1.1} (\text{syst})$
Σ_b^{*-}	$5835.0 \pm 0.6 (\text{stat}) \pm 1.8 (\text{syst})$	$6.4^{+2.2}_{-1.8} (\text{stat})^{+0.7}_{-1.1} (\text{syst})$

State	$\Delta M^{+-}, \text{ MeV}/c^2$
$\Sigma_b^+ - \Sigma_b^-$	$-4.2^{+1.1}_{-0.9} (\text{stat})^{+0.07}_{-0.09} (\text{syst})$
$\Sigma_b^{*+} - \Sigma_b^{*-}$	$-3.0 \pm 0.9 (\text{stat})^{+0.12}_{-0.13} (\text{syst})$

CDF Public Note 10286

Established existence of these states. First measurements of widths and mass splittings

ICHEP2010

*Spectroscopy of charmed $\Lambda_c(2595)$, $\Lambda_c(2625)$,
 $\Sigma_c(2455)$ and $\Sigma_c(2520)$ baryons*

ICHEP2010




Charm baryons – introduction

Λ_c^+ has $J^P = 1/2^+$ and is lightest c -baryon \rightarrow weak decay

Λ_c^* are orbital excitations of Λ_c^+

Σ_c are spin excitations of Λ_c^+ where light-diquark is spin-1

Isospin triplet decays strongly to $\Lambda_c^+\pi$






First CDF analysis of these modes – 5.2 fb⁻¹

Previous work mostly by CLEO




ICHEP2010

Charm baryons – reconstruction

Displaced-track triggered $\Lambda_c \rightarrow pK\pi$. Attach 1 (Σ_c) or 2 (Λ_c^*) pions. Two-step NN optimization: on Λ_c and on Σ_c/Λ_c^* using fit quality, lifetime p_T .





Neural Network



ICHEP2010

Charm baryons – Σ_c fits




Non-relativistic BW convoluted with detector resolution (from MC validated with $D^* \rightarrow D^0 \pi$ data).


Backgrounds from fake Λ_c , real Λ_c , + track, feeddown from $\Lambda_c(2625)$

ICHEP2010

Charm baryons – Λ_c^* fits



Mass-dependent width at threshold
PRD 67, 074033



Hard because $\Lambda_c^*(2595)$ is right above threshold for $\rightarrow \Sigma_c(2455)\pi$

Non-relativistic BW convoluted with detector resolution (from MC validated with $D^* \rightarrow D^0 \pi$ data).

Backgrounds from fake Λ_c , real Λ_c , + 2 tracks, $\Sigma_c(2455)$ +track

ICHEP2010

Charm baryons – results

Systematics: detector resolutions, mass scale (from $\psi(2S) \rightarrow J/\psi \pi \pi \pi$). Fit model. External input for reference masses.

	$m - m(\Lambda_c^+)[\text{MeV}/c^2]$	$\Gamma[\text{MeV}/c^2]$
$\Sigma_c(2455)^0$	167.28 ± 0.12 (167.30 ± 0.11)	1.65 ± 0.50 (2.2 ± 0.4)
$\Sigma_c(2455)^{++}$	167.44 ± 0.13 (167.56 ± 0.11)	2.34 ± 0.47 (2.23 ± 0.30)
$\Sigma_c(2520)^0$	232.88 ± 0.46 (231.6 ± 0.5)	12.51 ± 2.28 (16.1 ± 2.1)
$\Sigma_c(2520)^{++}$	230.73 ± 0.58 (231.9 ± 0.6)	15.03 ± 2.52 (14.9 ± 1.9)
$\Lambda_c(2595)^+$	<u>305.79 ± 0.24</u> (308.9 ± 0.6)	2.59 ± 0.56 ($3.6^{+2.0}_{-1.3}$)
$\Lambda_c(2625)^+$	341.65 ± 0.13 (341.7 ± 0.6)	$< 0.97(90\%CL)$ ($< 1.9(90\%CL)$)


- ❑ Comparable (Σ_c) or much better (Λ_c^*) than PDG. CDF Public Note 10260
- ❑ Difference in $\Lambda_c(2595)$ mass due to proper treatment of width at threshold. If not included get 3 MeV shift as seen in CLEO.
- ❑ Pion coupling constant determined from $\Lambda_c(2595)$: $h_2^2 = 0.36 \pm 0.08$

ICHEP2010

Observation of $Y(4140)$ exotic mesons

$Y(4140)$ – recap

2009: evidence of $J/\psi\phi$ structure at 4140 MeV in exclusive $B^+ \rightarrow J/\psi\phi K^+$





$M = 4143 \pm 2.9 \pm 1.2 \text{ MeV}$ (above open charm)

$\Gamma = 11.7^{+8.3}_{-5.0} \pm 3.7 \text{ MeV}$ (probably a strong decay)

Many exotic interpretations proposed. No signal seen by Belle which sets product of $\text{Br} < 6 \times 10^{-6}$ at 90%CL

$Y(4140) - 6\text{ fb}^{-1}\text{ analysis}$

Selection freezed to one used in 2009. Background shape changed.



S-wave BW convoluted with 1.7 MeV resolution + 3-body PS background

$$M = 4143.4^{+2.9}_{-3.0} \pm 0.6 \text{ MeV}$$


$$\Gamma = 15.3^{+10.4}_{-6.1} \pm 2.5 \text{ MeV}$$

Rate relative to $B^+ \rightarrow J/\psi\phi K^+$ is $(15 \pm 5)\%$

CDF Public Note 10244

ICHEP2010 $Y(4140) - 6 \text{ fb}^{-1}$ analysis (cont'd)

Prominent cluster of events at 4275 MeV




S-wave BW convoluted with 3.0 MeV resolution

$$M = 4274.4^{+8.4}_{-6.7} \pm 1.9 \text{ MeV} \quad \Gamma = 32.3^{+21.9}_{-15.3} \pm 7.6 \text{ MeV}$$

3σ significance

CDF Public Note 10244

Not done, yet



More than 10 fb^{-1} of physics-quality data on tape by end of 2011 (3 yrs extension under discussion)

Concluding remarks

Impressive Tevatron spectroscopy program. Data, highly efficient triggers, excellent tracking and high muon acceptance.

- ❑ World-leading mass determination of B^+ , B^0 , B_s^0 , Λ_b ground st. b-hadrons
- ❑ Pioneering studies of orbitally excited B^0 and B_s^0 mesons
- ❑ Till 2001, Λ_b was only known b -baryon. CDF/DØ filled-in gaps with observations of Σ_b , strange Ξ_b , doubly-strange Ω_b .
- ❑ Key contributions to exotic XYZ: new states, confirm B-factories, unique determination of quantum numbers, leading mass measurements.

Program ongoing and exploiting new opportunities.

3 recent results: improved Σ_b properties, entered realm of charmed baryons, yet another particle added to the zoo of exotic XYZ.

Sitting on a goldmine of ever increasing data. A few exciting years of competition with LHCb upcoming.

Tevatron Run II



www.fnal.gov